

New ways in testing post-Newtonian gravity in the Solar System scenario from planetary motion analysis

L. Iorio¹

Dipartimento di Fisica dell'Università di Bari, Via Amendola 173, 70126, Bari, Italy
e-mail: Lorenzo.Iorio@ba.infn.it

Received , 2004; accepted , 2004

Abstract. In this chapter we first calculate the post-Newtonian gravito-electric secular rate of the mean anomaly of a test particle freely orbiting a spherically symmetric central mass. Then, we propose a novel approach to suitably combine the presently available planetary ranging data to Mercury, Venus and Mars in order to determine, simultaneously and independently of each other, the Sun's quadrupole mass moment $J_{2\odot}$ and the post-Newtonian gravito-electric secular advances of the perihelion and the mean anomaly. This would also allow to obtain the PPN parameters γ and β independently. We propose to analyze the time series of three linear combinations of the observational residuals of the rates of the nodes $\dot{\Omega}$, the longitudes of perihelia $\dot{\pi}$ and mean anomalies $\dot{\mathcal{M}}$ of Mercury, Venus and Mars built up in order to account for the secular precessions induced by the solar oblateness and the post-Newtonian gravito-electric forces. According to the present-day EPM2000 and DE405 ephemerides accuracy, the obtainable precision would be of the order of 10^{-4} – 10^{-5} for the PPN parameters and, more interestingly, of 10^{-9} for $J_{2\odot}$. Second, we investigate the possibility of measuring the post-Newtonian Lense-Thirring secular effect induced by the gravito-magnetic field of the Sun on the planetary orbital motion by analyzing two linear combinations of the residuals of the nodes Ω of Mercury, Venus and Mars. The proposed combinations are, by construction, unaffected by the quadrupole mass moment J_2 of the Sun's gravitational potential. Moreover, they are insensitive also to the post-Newtonian gravito-electric field of the Sun because it affects only the perihelia ω and the mean anomalies $\mathcal{M}$. The obtainable observational accuracy in the proposed measurement would be of the order of 19% and 36%. However, it must be pointed out that a major source of systematic error could be represented by the residual mismodelled classical N -body secular precessions. The second proposed combination is insensitive just to such aliasing effect.

Key words. Relativity – Gravitation– Celestial Mechanics– Sun: fundamental parameters – Planets and satellites: general– Methods: miscellaneous

1. Introduction

In this chapter we wish to investigate the possibility of performing some tests of Newtonian and post-Newtonian gravity in the Solar System arena extending to it some approaches which, in recent years, have proven to be fruitful in certain Earth-artificial satellites systems. More precisely, we will deal with certain classical and post-Newtonian gravito-electromagnetic secular effects affecting the longitudes of the ascending nodes Ω , the longitudes of the perihelia π and the mean anomalies $\mathcal{M}$ of the Solar System's planets. The proposed tests could be done in a relatively simple and cheap way (with respect to new missions involving the construction and launch of one or more spacecrafts); moreover, they would also benefit of the improvements in the planetary ephemerides coming from future planned space missions like Messenger and BepiColombo. The obtainable accuracies are comparable with those of future proposed or planned expensive and complex interplanetary missions. In the case of the gravito-magnetic field, it would be the first attempt to measure its effect on the planetary orbital motion in the gravitational field of the Sun.

Historically, one of the first tests of the Einstein General Theory of Relativity (GTR) was the successful explanation of the anomalous secular gravito-electric perihelion advance of Mercury in the gravitational field of Sun (Einstein 1915). On the contrary, another prediction of the Einstein's theory of gravitation for the motion of test particles, i.e. the much smaller gravito-magnetic Lense-Thirring effect (Lense & Thirring 1918), is still awaiting for an undisputable

Table 1. Orbital parameters of Mercury, Venus and Mars (<http://nssdc.gsfc.nasa.gov/planetary/factsheet/>). For the Astronomical Unit (A.U.) we use the value 1 A.U.=149597870691 m of the DE405 (Standish 1998) and EPM2000 ephemerides (Pitjeva 2001a; 2001b). The angle ϵ refers to the inclination to the ecliptic.

Planet	a (A.U.)	ϵ ($^\circ$)	e
Mercury	0.38709893	7.00487	0.20563069
Venus	0.72333199	3.39471	0.00677323
Mars	1.52366231	1.85061	0.09341233

and direct observational check. As we will see, these post-Newtonian features of the planetary motion are strictly connected with the problem of the quadrupole mass moment $J_{2\odot}$ of Sun. In (Pireaux and Rozelot 2003) a theoretical range $J_{2\odot} = (2 \pm 0.4) \times 10^{-7}$ is admitted for the Sun's oblateness. For this topic and the interplay between this effect and the gravito-electric Einstein precession see the recent review (Pireaux & Rozelot 2003) and the references therein and (Ciufolini & Wheeler 1995). Basically, the point is the following. In regard to the secular orbital motions, to which we are interested here, the Sun's mass quadrupole moment induces classical effects which have, qualitatively, the same temporal signature of the relativistic ones. This means that they could corrupt the recovery of the genuine post-Newtonian features of motion because they could not be removed from the time series without removing the post-Newtonian signal of interest as well. All depends on the precision with which $J_{2\odot}$ is known: should the mismodelling in it induce classical residual precessions larger than the relativistic ones, there would be no hope to get a reliable test of relativistic gravity. As we will see later, the same problems could come from the N -body secular precessions. In this chapter we propose to disentangle such effects by measuring them in an independent way. More precisely, we propose to extend a certain approach used in Earth artificial satellite motion analysis to the interplanetary arena in order to single out just certain post-Newtonian (and Newtonian) orbital motion features independently of the Parameterized-Post-Newtonian (PPN) framework (Will 1993) which is usually employed in testing competing metric theories of gravity. Instead, the current approach consists of testing the post-Newtonian equations of motion as a whole in terms of the PPN parameters which are determined from multi-parameters fits together with other astrodynamical quantities.

In Table 1 the orbital parameters of the inner planets of the Solar System are reported.

2. The post-Newtonian effects

2.1. The gravito-electric effects

In the framework of the standard PPN formalism the post-Newtonian gravito-electric acceleration induced by the Schwarzschild-like part of the spacetime metric and experienced by a test body freely falling around a static, spherically symmetric central mass M is (Soffel 1989)

$$\mathbf{a}_{\text{GE}} = \frac{GM}{c^2 r^3} \left\{ \left[2(\gamma + \beta) \frac{GM}{r} - \gamma(\mathbf{v} \cdot \mathbf{v}) \right] \mathbf{r} + 2(1 + \gamma)(\mathbf{r} \cdot \mathbf{v})\mathbf{v} \right\}, \quad (1)$$

where G is the Newtonian gravitational constant, c is the speed of light in vacuum, $\mathbf{r}$ and $\mathbf{v}$ are the position and velocity vectors, respectively, of the test body, γ and β are the standard Eddington-Robertson-Schiff PPN parameters ($\gamma = \beta = 1$ in GTR; in this case eq.(1) reduces to the expression of (Mashhoon et al. 2001)). Note that r here is the standard isotropic radial coordinate, not to be confused with the Schwarzschild radial coordinate $r' = r[1 + GM/(2c^2 r)]^2$. In the usual orbital data reductions it is just r which is employed.

By considering it as a small perturbation to the Newtonian monopole acceleration it is possible to work out its effect on the orbital motion of a test body with the standard perturbative techniques. The secular rate of the argument of pericentre is given by the well known formula

$$\dot{\omega}_{\text{GE}} = \frac{3nGM}{c^2 a(1 - e^2)} \nu_{\text{GE}}, \quad (2)$$

where

$$\nu_{\text{GE}} = \frac{2 + 2\gamma - \beta}{3}, \quad (3)$$

a and e are the semimajor axis and the eccentricity, respectively, of the test particle's orbit and $n = \sqrt{GM/a^3}$ is the (unperturbed) Keplerian mean motion. The orbital period of an unperturbed, two-body Keplerian ellipse is $P = 2\pi/n$. Note that eq.(2) is an exact result valid to all order in e .

As we will show, also the mean anomaly $\mathcal{M}$ is affected by a post-Newtonian gravito-electric secular rate. The Gauss perturbative equation for $\mathcal{M}$ is given by

$$\frac{d\mathcal{M}}{dt} = n - \frac{2}{na} A_R \left(\frac{r}{a} \right) - \sqrt{1-e^2} \left(\frac{d\omega}{dt} + \cos i \frac{d\Omega}{dt} \right), \quad (4)$$

where Ω and i are the longitude of the ascending node and the inclination of the test particle's orbit to the equator of the central mass, respectively, and A_R is the radial component of the perturbing acceleration. In order to obtain the post-Newtonian gravito-electric secular rate of the mean anomaly one must consider the radial component of eq.(1), insert it in the second term of the right-hand-side of eq.(4), evaluate it on the unperturbed Keplerian ellipse, characterized by $r = a(1-e^2)/(1+e \cos f)$ and by the radial and along-track components of the velocity vector which are $v_R = nae \sin f / \sqrt{1-e^2}$, $v_T = na(1+e \cos f) / \sqrt{1-e^2}$, respectively (f is the orbiter's true anomaly), multiplying it by

$$\frac{dt}{P} = \frac{(1-e^2)^{3/2} df}{2\pi(1+e \cos f)^2}, \quad (5)$$

and integrating over one orbital revolution, i.e. from 0 to 2π . It turns out that the post-Newtonian gravito-electric secular rate of the mean anomaly is, to order $\mathcal{O}(e^2)$

$$\dot{\mathcal{M}}_{\text{GE}} \sim -\frac{nGM}{c^2 a} \left[(2+4\gamma+3\beta) \left(1 + \frac{e^2}{2} \right) + (2+\gamma) e^2 \right]. \quad (6)$$

We can define

$$\mu_{\text{GE}} = \frac{2+4\gamma+3\beta}{9}. \quad (7)$$

It turns out that the second term of eq.(6) induces for Mercury an additional shift of 1.789 arcseconds/century ($" \text{ cy}^{-1}$ in the following), while for the other planets it is of the order of 10^{-2} – $10^{-4} " \text{ cy}^{-1}$. Thus, for them the secular rate of the mean anomaly can be written as

$$\dot{\mathcal{M}}_{\text{GE}} \sim -\frac{9nGM}{c^2 a \sqrt{1-e^2}} \mu_{\text{GE}}. \quad (8)$$

However, as we will see later, the mean anomaly of Mercury will not be used in the combined residuals strategy outlined in the following. For Mercury, the post-Newtonian gravito-electric effect induced by the Sun on ω and $\mathcal{M}$, according to GTR, amounts to $42.980 " \text{ cy}^{-1}$ and $-127.949 " \text{ cy}^{-1}$, respectively. The most accurate estimate of the gravito-electric perihelion advance seems to be that obtained for Mercury by E.M. Standish in 2000 with the DE405 ephemerides (Standish 1998) and reported in (Pireaux and Rozelot 2003). He averaged the Mercury's perihelion evolution over two centuries by using the DE405 ephemerides with and without the post-Newtonian accelerations. Standish included in the force models also the solar oblateness with $J_{2\odot} = 2 \times 10^{-7}$, so that the residuals for Mercury accounted for the post-Newtonian effects only; the determined shift was $42.98 \pm 0.0023 " \text{ cy}^{-1}$. The same approach for the mean anomaly (Standish E M private communication) has yielded to $-130.003 \pm 0.0027 " \text{ cy}^{-1}$ (See also Table 4 later). Note that the quoted uncertainty do not come from direct observational errors. They depend on the fact that in the force models used in the numerical propagation many astrodynamical parameters occur (masses of planets, asteroids, etc.); their numerical values come from multiparameter fits of real data and, consequently, are affected by observational errors. Such numerical tests say nothing about if GTR is correct or not; they just give an idea of what would be the obtainable accuracy set up by our knowledge of the Solar System arena if the Einstein theory of gravitation would be true.

In regard to the possibility of constructing time series of planetary mean anomalies, it must be noted that a certain effort would be required. Indeed, so far this orbital elements has never been utilized, so that the partials, e.g., should be computed (Pitjeva, private communication 2004).

2.2. The gravito-magnetic Lense-Thirring effect

Another post-Newtonian secular precession which affects not only ω but also Ω is the Lense-Thirring effect (Lense & Thirring 1918) induced by the proper angular momentum J of the central body

$$\dot{\omega}_{\text{LT}} = \frac{-6GJ \cos i}{c^2 a^3 (1-e^2)^{3/2}} \mu_{\text{LT}}, \quad (9)$$

$$\dot{\Omega}_{\text{LT}} = \frac{2GJ}{c^2 a^3 (1-e^2)^{3/2}} \mu_{\text{LT}}, \quad (10)$$

where

$$\mu_{\text{LT}} = \frac{1 + \gamma}{2}. \quad (11)$$

By assuming for the Sun $J_{\odot} = 1.9 \times 10^{48} \text{ g cm}^2 \text{ s}^{-1}$ (Pijpers 2003), the gravito-magnetic effect on, e.g., Mercury's perihelion is of the order of $10^{-3}'' \text{ cy}^{-1}$ (De Sitter 1916). Such small value is at the edge of the present sensitivity ($0.002'' \text{ cy}^{-1}$) in determining the Mercury perihelion shift from the ephemerides (cfr. the results obtained by Standish). Moreover, it should be considered that, even if future improvements of the obtainable observational sensitivity in interplanetary ranging allowed to consider the possibility of measuring the Lense-Thirring perihelion advance of Mercury, the impact of the systematic errors due to the uncertainties in the solar oblateness would severely limit the realistic accuracy obtainable in such demanding measurement. Indeed, by assuming an uncertainty of $\sigma_{J_{2\odot}} = 0.4 \times 10^{-7}$, the error in the secular precession induced by the Sun's quadrupole moment would amount to 245% of the Lense-Thirring shift of π^{Merc} (see below for the definition of π). Another important source of systematic error in the measurement of such a tiny effect would be represented by the N -body classical secular precessions which are of the order of $10^2 - 10^3'' \text{ cy}^{-1}$ (see on the WEB <http://ssd.jpl.nasa.gov/elementplanets.html>). Indeed, their residual mismodelled part could severely bias the recovery of the Lense-Thirring effect. In this chapter we will propose a possible approach to overcome these problems. It must be noted that, if, on the other hand, we look at a Lense-Thirring test as a way to measure the Sun's angular momentum by assuming the validity of GTR, a measurement of the solar gravito-magnetic field would have a significance only if the obtainable accuracy was better than 10%; indeed, among other things, the present-day uncertainty in the Sun's angular momentum $J_{\odot}$, which could be measured from the Lense-Thirring precessions, is just of the order of 10% (Xu & Ni 1997) in various solar models (Paternó et al. 1996; Elsworth et al. 1995) or even less in the framework of asteroseismology (Pijpers 2003).

At present, the only performed attempts to explicitly extract the Lense-Thirring signature from the data of orbiting masses in the Solar System are due to Ciufolini and coworkers who analyzed the laser-ranged data of the orbits of the existing LAGEOS and LAGEOS II Earth artificial satellites (Ciufolini et al. 1998). A 20–30% precision level in measuring the terrestrial gravito-magnetic field is claimed, but other scientists judge these evaluations too optimistic and propose different error budget (Ries et al. 2003). In April 2004 the GP-B spacecraft has been launched. It will carry out a very complex and challenging mission which should be able to measure another gravito-magnetic effect, i.e. the precession of the spins of four superconducting gyroscopes (Schiff 1960) carried on board at a claimed accuracy of 1% or better (Everitt et al. 2001). Almost twenty years ago it was proposed to launch a third LAGEOS-like satellite—the LAGEOS III/LARES—and to analyze the time series of the sum of the residuals of the nodes of LAGEOS and LARES (Ciufolini 1986) or some other combinations of residuals of the nodes and the perigees of LARES and both the existing LAGEOS satellites (Iorio et al. 2002a). The obtainable accuracy would probably be of the order of 1% too. Mainly funding problems have prevented, up to now, from implementing such relatively easy and cheap mission. Recently, the possibility of measuring the Lense-Thirring precessions by means of the Relativity-dedicated OPTIS spacecraft, which could be launched in the same orbital configuration of LARES, has been considered (Iorio et al., 2004). The recently proposed LATOR (Turyshev et al. 2004) and ASTROD (Ni et al. 2004) missions would be sensitive to the gravito-magnetic part of the bending of light rays and time delay in the gravitational field of the Sun. Finally, it must be noted that, according to K. Nordvedt Jr. (Nordvedt 2003), the multidecadal analysis of the Moon's motion with the Lunar Laser Ranging (LLR) technique strongly supports the existence of the gravito-magnetic force¹ as predicted by GTR, although in an indirect way. “It often has been claimed that the presence of gravitomagnetism within the total gravitational interaction has not been experimentally confirmed and measured. Indeed, different experiments have been under development to explicitly observe the effects of this historically interesting prediction of general relativity. But this gravito-magnetic acceleration already plays a large role in producing the final shape of the lunar orbit, albeit in conjunction with the rest of the total equation of motion; the precision fit of the LLR data indicates that gravitomagnetism's presence and specific strength in the equation of motion can hardly be in doubt. [...] It would be impossible to understand this fit of the LLR data without the participation of the gravito-magnetic interaction in the underlying model, and with strength very close to that provided by general relativity, $\gamma = 1$ ”.

3. The solar oblateness

The solar quadrupole mass moment $J_{2\odot}$ is an important astrophysical parameter whose precise knowledge could yield many information about the inner structure and dynamics of our star. A reliable evaluation of $J_{2\odot}$ still faces some controversy: on one side, the theoretical values strongly depend on the solar model used, whereas accurate

¹ According to Nordvedt (2003), the Earth–Moon range is affected by long-periodic harmonic perturbations of gravito-magnetic origin whose amplitudes are of the order of 5 m and the periods are monthly and semi-monthly. The amplitudes of the lunar motion at both these periods are determined to better than half a centimeter precision in the total orbital fit to the LLR data.

Table 2. Post-Newtonian gravito-electric precessions and coefficients of Newtonian precessions of the node, the perihelion and the mean anomaly for Mercury, Venus and Mars in $'' \text{cy}^{-1}$. The values $\dot{\pi}_{\text{GE}}$, and $\dot{\mathcal{M}}_{\text{GE}}$ are calculated with GTR. The coefficients $\dot{\Omega}_{.2}$, $\dot{\pi}_{.2}$ and $\dot{\mathcal{M}}_{.2}$ are $\partial(\dot{\Omega}_{J_{2\odot}})/\partial(J_{2\odot})$, $\partial(\dot{\pi}_{J_{2\odot}})/\partial(J_{2\odot})$ and $\partial(\dot{\mathcal{M}}_{J_{2\odot}})/\partial(J_{2\odot})$, respectively. For Ω and ω they can be found in (Iorio 2003). In order to have the precessions they must be multiplied by $J_{2\odot}$. Note that the result for the mean anomaly of Mercury accounts for the correction of order $\mathcal{O}(e^2)$ which, instead, can be neglected for the other planets.

Planet	$\dot{\Omega}_{\text{GE}}$	$\dot{\pi}_{\text{GE}}$	$\dot{\mathcal{M}}_{\text{GE}}$	$\dot{\Omega}_{.2}$	$\dot{\pi}_{.2}$	$\dot{\mathcal{M}}_{.2}$
Mercury	0	42.981	-127.949	-126878.626	126404.437	123703.132
Venus	0	8.624	-25.874	-13068.273	13056.803	13056.504
Mars	0	1.351	-4.035	-980.609	980.353	976.067

measurements are very difficult to obtain from observations. For all this matter see the recent review (Pireaux & Rozelot 2003) and (Rozelot et al. 2004). From an observational point of view, $J_{2\odot}$ is not directly accessible. In this context a dynamical determination of $J_{2\odot}$, analyzing, e.g., the orbits of the inner planets of the Solar System, is interesting because it might be compared with those derived from solar model dependent values of the oblateness. However, it is not simple to reach this goal because of the interplay between the effects of the solar quadrupole moment with those induced by the post-Newtonian gravito-electromagnetic forces. Instead of the trajectory of planets, it would be possible to infer $J_{2\odot}$ from accurate tracking of some drag-free spacecraft orbiting within a few radii of the solar center. This will be the approach followed by, e.g. the BepiColombo mission (see Section 4.2). Alternatively, the Sun's quadrupole mass moment can be inferred from in-orbit measurement of solar properties, like the SOHO-*MDI* space-based observations (Armstrong & Kuhn 1999), or from Earth-based observations like those realized, e.g., with the scanning heliometer of the Pic du Midi Observatory (Rozelot et al. 2004).

3.1. The classical precessions induced by the solar oblateness

For a given planet of the Solar System orbiting the Sun, apart from the classical effects induced by the precession of the equinoxes and by the other planets and major asteroids which are routinely accounted for in the ephemerides computations (Pitjeva 2001a; 2001b), the oblateness of Sun induces also secular precessions on Ω , $\pi = \omega + \Omega \cos i$ and $\mathcal{M}$ given by

$$\dot{\Omega}_{J_{2\odot}} = -\frac{3}{2} \frac{nJ_{2\odot}}{(1-e^2)^2} \left(\frac{R_{\odot}}{a}\right)^2 \cos i, \quad (12)$$

$$\dot{\pi}_{J_{2\odot}} = -\frac{3}{2} \frac{nJ_{2\odot}}{(1-e^2)^2} \left(\frac{R_{\odot}}{a}\right)^2 \left(\frac{3}{2} \sin^2 i - 1\right), \quad (13)$$

$$\dot{\mathcal{M}}_{J_{2\odot}} = \frac{3}{4} \frac{nJ_{2\odot}}{(1-e^2)^{3/2}} \left(\frac{R_{\odot}}{a}\right)^2 (3 \cos^2 i - 1), \quad (14)$$

The explicit expressions for the secular precessions of Ω and ω up to degree $\ell = 20$ can be found in (Iorio 2003). For Mercury² the Newtonian precessions due to Sun oblateness, with $J_{2\odot} = 2 \times 10^{-7}$, are of the order of $10^{-2} '' \text{cy}^{-1}$ for Ω , π and $\mathcal{M}$.

4. The interplay between the solar oblateness and the post-Newtonian precessions

In Table 2 the relevant parameters of the classical and post-Newtonian gravito-electric secular precessions of the nodes, the perihelia and the mean anomalies of Mercury, Venus and Mars are reported. In all the relativistic tests performed up to now by analyzing the perihelia advances only of the inner planets of the Solar System with the radar ranging technique (Shapiro et al. 1972; 1976; Shapiro 1990) it has been impossible to disentangle the genuine post-Newtonian gravito-electric contribution of eq.(2) from the Newtonian precession of eq.(13). Indeed, the observational residuals of $\dot{\pi}$ for a single planet, built up by suitably switching off the post-Newtonian $\mathcal{O}(c^{-2})$ terms and the oblateness of

² As pointed out in (Milani et al., 2002), the angle i refers to the inclination between the planet's orbital plane and the fixed reference plane of the celestial reference frame; it is not the angle ϵ between the planet's orbital plane and the ecliptic. It turns out that $i \sim \epsilon/2$. For Mercury $\epsilon = 7.00487^\circ$.

Table 3. Post-Newtonian gravito-magnetic precessions and coefficients of Newtonian precessions of the node for Mercury, Venus and Mars in $'' \text{cy}^{-1}$. The values $\dot{\Omega}_{\text{LT}}$ are calculated with GTR. The coefficients $\dot{\Omega}_{\ell}$, are $\partial(\dot{\Omega}_{J_{\ell\odot}})/\partial(J_{\ell\odot})$. They can be found, up to degree $\ell = 20$, in (Iorio 2003). In order to have the precessions they must be multiplied by $J_{\ell\odot}$.

Planet	$\dot{\Omega}_{\text{LT}}$	$\dot{\Omega}_{.2}$	$\dot{\Omega}_{.4}$	$\dot{\Omega}_{\text{N-body}}$
Mercury	0.001008	-126878.626476	52.774935	-446.30
Venus	0.000144	-13068.273031	1.349709	-996.89
Mars	0.000015	-980.609460	0.023554	-1020.19

Sun in the force models of the equations of motion in the orbital processors softwares, account entirely for the post-Newtonian and the solar oblateness³ effects. This is a unsatisfactory situation, both if we are interested in testing post-Newtonian gravity and if we want to obtain a dynamical, model-independent measurement of $J_{2\odot}$. Indeed, it is, of course, impossible to constraint both the effects if only one perihelion rate is examined one at a time: in recovering one of the two effects we are forced to consider the other one as if it was known. Since the post-Newtonian gravito-electric effect is three orders of magnitude larger than that induced by solar oblateness, a determination of the latter by assuming the validity of GTR would be affected by a non negligible systematic error induced by the precision to which the post-Newtonian pericentre advance is known from other (more or less indirect and more or less biased by other aliasing effects) tests (Lunar Laser Ranging, binary pulsars periastron advance⁴). The inverse situation is more favorable: indeed, if we are interested in the post-Newtonian gravito-electric effect the relative systematic error induced on its measurement by the precession due to the solar oblateness amounts to 5×10^{-4} even by assuming for the latter effect a 100% uncertainty⁵.

4.1. The present-day approach to test post-Newtonian gravity

At this point it may be interesting to clarify what is the current approach in testing post-Newtonian gravity from planetary data analysis followed by, e.g., the Jet Propulsion Laboratory (JPL). In the interplay between the real data and the equations of motions, which include also the post-Newtonian accelerations expressed in terms of the various PPN parameters, a set of astrodynamical parameters, among which there are also γ and β , are simultaneously and straightforwardly fitted and adjusted and a correlation matrix is also released. This means that the post-Newtonian equations of motion are globally tested as a whole in terms of, among other parameters, γ and β ; no attention is paid to this or that particular feature of the post-Newtonian accelerations. The point is that the standard PPN formalism refers to the alternative theories of gravitation which are metric, i.e. based on a symmetric spacetime metric. But it is not proven that an alternative theory of gravitation must necessarily be a metric one. Moreover, even in the framework of the metric alternative theories, the PPN formalism based on 10 parameters is not sufficient to describe every conceivable metric theory of gravitation at the post-Newtonian order; it only describes those theories with a particularly simple post-Newtonian limit. One would, in principle, need an infinite set of new parameters to add to the standard ten parameter PPN formalism in order to describe the post-Newtonian approximation of any a priori conceivable metric theory of gravity (Ciufolini 1991; Ciufolini & Wheeler 1995).

4.2. The possibilities opened by the future missions

Concerning the possibility of disentangle the effects of the solar oblateness from those of the post-Newtonian gravito-electric force, it is stated that the future space mission BepiColombo⁶ of the European Space Agency (ESA) will

³ Also if the solar oblateness is included in the force models, the related uncertainty induces a corresponding systematic error in the recovered post-Newtonian effect. Fortunately, it is small; by assuming $\sigma_{J_{2\odot}} = 0.4 \times 10^{-7}$, it amounts to 0.01%.

⁴ Note that the binary pulsars periastron measurement should not be considered as a test of relativistic gravity because the masses of the binary system are not known (Stairs et al. 1998). The Lunar Laser Ranging measurements do not allow to single out uniquely the gravitoelctric pericentre advance from the other post-Newtonian features of motion of the Earth-Moon system (Nordvedt 2001). Recently, it has been proposed to measure the relativistic gravito-electric perigee advance of the terrestrial LAGEOS II satellite (Iorio et al. 2002b; Lucchesi 2003), but, up to now, the experiment has not yet been performed.

⁵ Note that the impossibility of disentangle the gravito-electric and Lense-Thirring effects would not seriously affect the recovery of the gravitoelctric precession: indeed, the bias induced by the gravito-magnetic effect on the gravito-electric shift amounts to 0.004% only for Mercury.

⁶ See on the WEB <http://astro.estec.esa.nl/BepiColombo/>. Present ESA plans are for a launch in 2010-2012.

Table 4. Present-day accuracy in determining the node, perihelion and mean anomaly secular rates of Mercury, Venus and Mars according to DE405 (Standish 1998) and EPM2000 (Pitjeva 2001a; 2001b) ephemerides. The figures, in " cy^{-1} , represent the formal, statistical errors. Realistic errors should be 10 times larger, at least. While the results by Standish come from the mathematical propagation of the nodes, the perihelia and the mean anomalies (Standish, E. M. private communication) evolution with and without post-Newtonian terms (with $\gamma = \beta = 1$) and their average over a time span of two centuries, the results by Pitjeva are based on real data. The figure for Mercury has been obtained in 2001, while the other ones have been determined subsequently (Pitjeva, personal communication 2004).

Planet	$\sigma_{\dot{\Omega}_{\text{calc}}}(DE405)$	$\sigma_{\dot{\pi}_{\text{calc}}}(DE405)$	$\sigma_{\dot{\mathcal{M}}_{\text{calc}}}(DE405)$	$\sigma_{\dot{\pi}_{\text{obs}}}(EPM2000)$
Mercury	0.000182	0.0023	0.0027	0.0086
Venus	0.000006	0.0414	0.0414	0.1037
Mars	0.000001	0.0014	0.0014	0.0001

provide us, among other things, with a dynamical, model-independent and relativity-independent measurement of $J_{2\odot}$ by measuring with high precision the nodal motion of Mercury (Milani et al. 2002; Pireaux & Rozelet 2003) which is not affected by the post-Newtonian gravito-electric force. The claimed accuracy would amount to $\sigma_{J_{2\odot}} = 2 \times 10^{-9}$ (Milani et al. 2002). However, such evaluation refers to the formal, statistical obtainable uncertainty only. Indeed, the residuals of the Mercury's node would account, to a certain level of accuracy, for the Lense-Thirring precession as well. By considering such effect as totally unmodelled in the force models, its impact on the measurement of $J_{2\odot}$ would induce a 8×10^{-9} systematic error. The formal, statistical accuracy for γ and β is evaluated to be of the order of 2×10^{-6} (Milani et al. 2002). Also the ESA astrometric mission GAIA⁷ should measure, among other things, the solar quadrupole mass moment by analyzing the longitudes of the ascending nodes of many minor bodies of the Solar System. The obtainable accuracy for γ is of the order of 10^{-5} - 10^{-7} (Vecchiato et al. 2003). The ASTROD mission should be able to measure $J_{2\odot}$ with a claimed accuracy of the order of 10^{-8} or, perhaps, $10^{-9} - 10^{-10}$ (Ni et al. 2004). The claimed obtainable accuracy for the PPN parameters is 4.6×10^{-7} for γ and 4×10^{-7} for β . Further improvements may push these limits down to $10^{-8} - 10^{-9}$. The recently proposed LATOR mission should be able to measure, among other things, γ to a 10^{-8} accuracy level and $J_{2\odot}$ to a 10^{-8} level (Turyshev et al. 2004). However, LATOR, whose goal is an extremely accurate measurement of the light deflection at the Sun's limb, should not allow to disentangle the various post-Newtonian contributions (first and second order gravito-electric contributions, first order gravito-magnetic term, solar oblateness effect) to it. The solar orbit relativity test SORT (Melliti et al. 2002), which would combine a time-delay experiment with a light deflection test, should allow to reach a 10^{-6} accuracy in measuring the PPN parameters

5. The observational accuracy in planetary radar ranging

Concerning the present-day accuracy of the planetary radar ranging, the radar itself is accurate well below the 100 m level (Standish 2002). The problem, however, comes from the fact that the surfaces of the planets have large topographical variations. They are modeled in different ways. For Mercury, spherical harmonics and some closure analysis (comparing values when two different measurements reflect off from the same spot on the surface) have been done (Anderson et al. 1996) in DE405. For Venus, a topographical model, which comes from (Pettengill et al. 1980), has been used. For Mars, closure points can be used. Closure points are pairs of days during which the observed points on the surface of Mars are nearly identical with respect to their longitudes and latitudes on Mars. Since the same topographical features are observed during each of the two days, the uncertainty introduced by the topography may be eliminated by subtracting the residuals of one day from the corresponding ones of the other day. The remaining difference is then due to only the ephemeris drift between the two days. The closure points for Mars have a priori uncertainties of about 100 m or less when the points are within 0.2 degrees of each other on the martian surface. Of course, for Mars, there is also the spacecraft ranging - far more accurate than the radar: Viking Landers (1976-82), 10 m; Mars Global Surveyor and Odyssey (1999-2003), 2-3 m. However, correction for Mars topography is possible not only by using closure points (in this method no all observations may be used), but with help of modern hypsometric maps and by the representations of the global topography with an expansion of spherical functions (Pitjeva 2001b).

⁷ See on the WEB <http://astro.estec.esa.nl/GAIA/>. Present ESA plans are for a launch in mid-2010.

6. The proposed approach for disentangle the solar oblateness and the post-Newtonian effects

In regard to the possibility of measuring explicitly some post-Newtonian effects from the analysis of the orbital motion of proof masses in the gravitational field of a real (rotating) astronomical mass like the Earth, it must be pointed out that the main problems come from the aliasing effects induced by a host of classical orbital perturbations, of gravitational and non-gravitational origin, which unavoidably affect the motion of the probes along with GTR. In particular, the even zonal harmonics J_ℓ of the multipolar expansion of the gravitational potential of the central mass induce secular classical precessions which, in many cases, tend to alias the gravito-electromagnetic ones of interest. Moreover, also the non-gravitational perturbations, to which the perigees of the LAGEOS-like satellites are particularly sensitive, are another important source of bias. The approach proposed by Ciufolini (1996) and Iorio (Iorio 2002; Iorio & Morea 2004) in the performed or proposed tests with LAGEOS and LAGEOS II consists of suitably designing linear combinations $\sum u_i \delta \dot{\Omega}_{\text{obs}}^i + \sum v_j \delta \dot{\omega}_{\text{obs}}^j$ of orbital residuals which are able to reduce the impact both of the even zonal harmonics of the gravitational field of the central mass and of the non-gravitational perturbations. In general, the coefficients u_i and v_j which weigh the various orbital elements in the combinations are a compromise between these two distinct needs.

6.1. The gravito-electric precessions and the solar oblateness

The main idea is to extend this approach to the Sun-planets system. It must be considered that, in this case, the non-gravitational perturbations do not play any role⁸, while the N -body classical perturbations of gravitational origin are important. Let us start with the gravito-electric precessions writing down the following equations⁹

$$\begin{cases} \delta \dot{\Omega}_{\text{obs}}^{\text{Merc}} = \dot{\Omega}_{.2}^{\text{Merc}} J_{2\odot} + \dot{\Omega}_{\text{N-body}}^{\text{Merc}} + \dot{\Omega}_{\text{LT}}^{\text{Merc}} \mu_{\text{LT}}, \\ \delta \dot{\Omega}_{\text{obs}}^{\text{Venus}} = \dot{\Omega}_{.2}^{\text{Venus}} J_{2\odot} + \dot{\Omega}_{\text{N-body}}^{\text{Venus}} + \dot{\Omega}_{\text{LT}}^{\text{Venus}} \mu_{\text{LT}}, \\ \delta \dot{\Omega}_{\text{obs}}^{\text{Mars}} = \dot{\Omega}_{.2}^{\text{Mars}} J_{2\odot} + \dot{\Omega}_{\text{N-body}}^{\text{Mars}} + \dot{\Omega}_{\text{LT}}^{\text{Mars}} \mu_{\text{LT}}, \end{cases} \quad (15)$$

$$\begin{cases} \delta \dot{\mathcal{M}}_{\text{obs}}^{\text{Mars}} = \dot{\mathcal{M}}_{.2}^{\text{Mars}} J_{2\odot} + \dot{\mathcal{M}}_{\text{GE}}^{\text{Mars}} \mu_{\text{GE}}, \\ \delta \dot{\mathcal{M}}_{\text{obs}}^{\text{Venus}} = \dot{\mathcal{M}}_{.2}^{\text{Venus}} J_{2\odot} + \dot{\mathcal{M}}_{\text{GE}}^{\text{Venus}} \mu_{\text{GE}}, \end{cases} \quad (16)$$

$$\begin{cases} \delta \dot{\pi}_{\text{obs}}^{\text{Mars}} = \dot{\pi}_{.2}^{\text{Mars}} J_{2\odot} + \dot{\pi}_{\text{GE}}^{\text{Mars}} \nu_{\text{GE}}, \\ \delta \dot{\pi}_{\text{obs}}^{\text{Merc}} = \dot{\pi}_{.2}^{\text{Merc}} J_{2\odot} + \dot{\pi}_{\text{GE}}^{\text{Merc}} \nu_{\text{GE}}, \end{cases} \quad (17)$$

where $\delta \dot{\Omega}_{\text{obs}}^{\text{Planet}}$, $\delta \dot{\pi}_{\text{obs}}^{\text{Planet}}$ and $\delta \dot{\mathcal{M}}_{\text{obs}}^{\text{Planet}}$ are the observational residuals of the rates of the nodes, the longitudes of the perihelia and the mean anomalies of Mercury, Venus and Mars. It is intended that all kind of data (optical and radio) would be used. The residuals¹⁰ should be built up by purposely switching off the solar quadrupole moment and the post-Newtonian gravito-electric accelerations in the force models (or leaving in them some default values to be subsequently adjusted according to the present strategy) of the orbital processors. Then, the so obtained observational residuals would entirely (or partly, if some default values are left in the force models) adsorb just the investigated secular effects and other post-Newtonian short-periodic features, i.e. not averaged over one orbital revolution of the planet under consideration. In respect to the latter point, it should be noted that the residuals of the mean anomalies

⁸ Indeed, they are proportional to the area-to-mass ratio S/M of the orbiting probe. The area-to-mass ratio falls off as $1/r$ where r is the radius of the probe-a planet, in this case-assumed spherical.

⁹ Note that in the right-hand-sides of eqs.(16)-(17) also the mismodelled parts of the classical N -body precessions should have been included. Since they are small with respect to the gravito-electric effects of interest, as we will show later, we can neglect them in the calculations for setting up our combinations. This, of course, does not mean that the residuals do not account also for them.

¹⁰ Note that here we use expressions like ‘observational residuals of a Keplerian orbital element’, strictly speaking, in an improper sense. The true observables, for which it is possible to build up residuals, are ranges, range rates and angles. The Keplerian orbital elements are not directly observable: they can only be calculated from an orbit which has been either numerically propagated or fitted to real observational data (the observed orbit) starting from given initial conditions. Here with ‘time series of the residuals of a Keplerian element’ we mean the difference between the time series of that Keplerian element related to an observed orbital arc and the time series of that Keplerian element numerically propagated on the same orbital arc (equal initial conditions) with the given classical or post-Newtonian parameter of interest switched off in the force models.

would account, e.g., also for the indirect effects on the mean motions n through the perturbations in the semimajor axes a

$$\Delta n = -\frac{3}{2}\sqrt{\frac{GM}{a^5}}\Delta a. \quad (18)$$

There are no secular perturbations induced on a by the other planets. If the classical short-periodic effects on a would be of relatively no importance because they would be included in the force models at the best of their accuracy, this is not the case for the post-Newtonian ones. The gravito-electric field induces no secular variations on a , as the classical planetary perturbations. The short-term shift on a can be calculated from eq.(1) and the Gauss equation for the perturbed rate of semimajor axis

$$\frac{da}{dt} = \frac{2}{n\sqrt{1-e^2}} [A_R e \sin f + A_T (1 + e \cos f)], \quad (19)$$

where A_T is the along-track component of the perturbing acceleration. It amounts to

$$\Delta a_{GE} = \frac{GMe}{c^2(1-e^2)^2} [14(\cos f_0 - \cos f) + 10e(\cos^2 f - \cos^2 f_0)] + \mathcal{O}(e^3). \quad (20)$$

For Venus their amplitudes are of the order of 100 m; for Mars they amount to 2 km; eq.(18) would yield periodic variations whose amplitudes would be of the order of 0.2-1 '' cy⁻¹ for Venus and Mars, respectively. However, such harmonic signatures could be fitted and removed from the data over sufficiently long time spans. Indeed, since we are interested in the gravito-electric secular trends on π and $\mathcal{M}$ it should be possible, in principle, to construct the residuals by using orbital arcs longer than the sidereal revolution periods of the planets to be used. Then, all the high-frequency perturbations would not affect their time series which should, instead, be characterized by the secular parts of those Newtonian and post-Newtonian features present in the real data but (partly) absent in the force models of the equations of motion in the orbital processors. However, in regard to the possibility of constructing accurate time series of observational residuals many years long the following observations must be kept in mind (Standish 2002). The planetary motions are perturbed by the presence of many asteroids whose masses are quite poorly known. Furthermore, it is not possible to solve for the asteroid masses, other than for the biggest few, because there are too many of them for the data to support such an effort. As a result, the ephemerides of the inner planets, especially Mars, will deteriorate over time; the ephemerides have uncertainties at the 1-2 km level over the span of the observations and growing at the rate of a few km/decade outside that span. On the other hand, it must also be noted that the sidereal orbital periods of Mercury and Mars amount to 87.969 and 686.980 days, respectively.

We can consider eqs.(15)-(17) as three systems of algebraic linear equations in the three unknowns¹¹ $J_{2\odot}$, μ_{GE} and ν_{GE} . Their solutions can be written as

$$\begin{cases} \delta\dot{\Omega}_{\text{obs}}^{\text{Merc}} + c_1\delta\dot{\Omega}_{\text{obs}}^{\text{Venus}} + c_2\delta\dot{\Omega}_{\text{obs}}^{\text{Mars}} = J_{2\odot} \left(\dot{\Omega}_{.2}^{\text{Merc}} + c_1\dot{\Omega}_{.2}^{\text{Venus}} + c_2\dot{\Omega}_{.2}^{\text{Mars}} \right), \\ \delta\dot{\mathcal{M}}_{\text{obs}}^{\text{Mars}} + c_1'\dot{\mathcal{M}}_{\text{obs}}^{\text{Venus}} = \mu_{GE} \left(\dot{\mathcal{M}}_{\text{GE}}^{\text{Mars}} + c_1'\dot{\mathcal{M}}_{\text{GE}}^{\text{Venus}} \right), \\ \delta\dot{\pi}_{\text{obs}}^{\text{Mars}} + c_1''\dot{\pi}_{\text{obs}}^{\text{Merc}} = \nu_{GE} \left(\dot{\pi}_{\text{GE}}^{\text{Mars}} + c_1''\dot{\pi}_{\text{GE}}^{\text{Merc}} \right), \end{cases} \quad (21)$$

where

$$\begin{cases} c_1 = \frac{\dot{\Omega}_{\text{LT}}^{\text{Mars}}\dot{\Omega}_{\text{N-body}}^{\text{Merc}} - \dot{\Omega}_{\text{LT}}^{\text{Merc}}\dot{\Omega}_{\text{N-body}}^{\text{Mars}}}{\dot{\Omega}_{\text{LT}}^{\text{Venus}}\dot{\Omega}_{\text{N-body}}^{\text{Mars}} - \dot{\Omega}_{\text{LT}}^{\text{Mars}}\dot{\Omega}_{\text{N-body}}^{\text{Venus}}} = -7.73247, \quad c_2 = \frac{\dot{\Omega}_{\text{LT}}^{\text{Merc}}\dot{\Omega}_{\text{N-body}}^{\text{Venus}} - \dot{\Omega}_{\text{LT}}^{\text{Venus}}\dot{\Omega}_{\text{N-body}}^{\text{Merc}}}{\dot{\Omega}_{\text{LT}}^{\text{Venus}}\dot{\Omega}_{\text{N-body}}^{\text{Mars}} - \dot{\Omega}_{\text{LT}}^{\text{Mars}}\dot{\Omega}_{\text{N-body}}^{\text{Venus}}} = 7.11840 \\ c_1' = -\dot{\mathcal{M}}_{.2}^{\text{Mars}}/\dot{\mathcal{M}}_{.2}^{\text{Venus}} = -0.07475, \\ c_1'' = -\dot{\pi}_{.2}^{\text{Mars}}/\dot{\pi}_{.2}^{\text{Merc}} = -0.00775, \end{cases} \quad (22)$$

and

$$\begin{cases} \dot{\Omega}_{.2}^{\text{Merc}} + c_1\dot{\Omega}_{.2}^{\text{Venus}} + c_2\dot{\Omega}_{.2}^{\text{Mars}} = -32808.8816'' \text{ cy}^{-1}, \\ \dot{\mathcal{M}}_{\text{GE}}^{\text{Mars}} + c_1'\dot{\mathcal{M}}_{\text{GE}}^{\text{Venus}} = -2.1007'' \text{ cy}^{-1}, \\ \dot{\pi}_{\text{GE}}^{\text{Mars}} + c_1''\dot{\pi}_{\text{GE}}^{\text{Merc}} = 1.0176'' \text{ cy}^{-1} \end{cases} \quad (23)$$

¹¹ This approach is analogous to that employed in the LAGEOS-LAGEOS II Lense-Thirring experiment in the gravitational field of Earth (Ciufolini 1996; Ciufolini et al. 1998)

The first equation of eq.(21) comes from eq.(15) solved for $J_{2\odot}$; it allows to obtain $J_{2\odot}$ independently of the post-Newtonian Lense-Thirring and classical N -body secular precessions which would represent the major sources of systematic errors. The second equation of eq.(21) comes from eq.(16) solved for μ_{GE} ; it cancels out the secular precessions due to $J_{2\odot}$. The third equation in eq.(21) comes from eq.(17) solved for ν_{GE} ; it cancels out the secular precessions due to $J_{2\odot}$. In regard to the impact of the residual mismodelled classical N -body precessions on the second and the third combinations of eq.(21), they should not induce a systematic error larger than the observational one (see below) because the expected values of μ_{GE} and ν_{GE} are of the order of unity¹², contrary to $J_{2\odot}$ which should be of the order of 10^{-7} . The adimensional parameters $J_{2\odot}$, μ_{GE} and ν_{GE} are estimated by fitting the time series of the left-hand-sides of eq.(21) with straight lines, measuring their slopes, in $'' \text{cy}^{-1}$, and, then, by dividing them by the quantities of eq.(23) which have the dimensions of $'' \text{cy}^{-1}$. Note that the solar quadrupole mass moment would not be affected by the indirect effects on n because only the nodes would be used in its determination. Finally, from the so obtained values of μ_{GE} and ν_{GE} , which are 1 in GTR and 0 in Newtonian mechanics, it is possible to measure γ and β independently of the solar oblateness and also of each other as

$$\begin{cases} \gamma = \frac{9}{10}(\mu_{\text{GE}} + \nu_{\text{GE}}) - \frac{4}{5}, \\ \beta = \frac{9}{5}\mu_{\text{GE}} - \frac{6}{5}\nu_{\text{GE}} + \frac{2}{5}. \end{cases} \quad (24)$$

According to the results of Table 4 it is possible to yield an estimate of the (formal) uncertainty in $J_{2\odot}$, μ_{GE} and ν_{GE} as

$$\begin{cases} \sigma_{J_{2\odot}} \sim 4.5 \times 10^{-9}, \\ \sigma_{\mu_{\text{GE}}} \sim 6 \times 10^{-5}, \\ \sigma_{\nu_{\text{GE}}} \sim 3 \times 10^{-5}. \end{cases} \quad (25)$$

The evaluation for $\sigma_{\mu_{\text{GE}}}$ accounts also for the fact that eq.(18) and the values of σ_a of Table VI of (Pitjeva 2001a) yield a (formal) uncertainty of 3×10^{-5} . The uncertainties in γ and β would amount to

$$\begin{cases} \sigma_{\gamma} \sim 6 \times 10^{-5}, \\ \sigma_{\beta} \sim 1 \times 10^{-4}. \end{cases} \quad (26)$$

However, discretion is advised on evaluating the reliability of these results because they refer to the formal, standard statistical errors; realistic errors may be also one order of magnitude larger.

For the most recent determinations of γ and β we have that, according to the frequency shift of radio photons to and from the Cassini spacecraft, $\sigma_{\gamma} = 2 \times 10^{-5}$ (Bertotti et al. 2003). This result, combined with that for the Strong Equivalence Principle violating parameter $\sigma_{\eta} = 4.5 \times 10^{-4}$ from the most recent analysis of LLR data, yields $\sigma_{\beta} = 1 \times 10^{-4}$ (Turyshev et al. 2003).

6.2. The gravito-magnetic precessions

Let us, now, turn our attention to the Lense-Thirring effect and try to set up some combinations with the nodes of the inner planets which cancel out the impact of the Sun's oblateness and of the N -body precessions.

6.2.1. A $J_{2\odot} - J_{4\odot}$ free combination

Let us write down the expressions of the residuals of the nodes of Mercury, Venus and Mars explicitly in terms of the mismodelled secular precessions induced by the quadrupolar and octupolar (Rozelet et al. 2004) mass moments of the

¹² Let us quantitatively discuss this point. By using the results for the observed centennial rates $\dot{\Omega}$ and $\dot{\omega}$ released at http://ssd.jpl.nasa.gov/elem_planets.html in, say, the left-hand-side of the third combination of eq.(21) it is possible to obtain for it a nominal N -body shift of $539.6036'' \text{cy}^{-1}$. The uncertainty in the N -body precessions lies mainly in the Gm of the perturbing planets, among which Jupiter plays the major role. Now, the relative uncertainty in Jupiter's Gm is of the order of 10^{-8} (Jacobson 2003); then, a reasonable estimate of the order of magnitude of the mismodelled part of the N -body shift should be $1 \times 10^{-5}'' \text{cy}^{-1}$. This figure must be divided by $1.0176'' \text{cy}^{-1}$ yielding a relative error in ν_{GE} of the order of 10^{-5} . Cfr. with the third equations of eq.(25).

Sun and of the Lense-Thirring secular precessions, assumed as a totally unmodelled feature. It is accounted for by a scaling parameter μ_{LT} which is zero in Newtonian mechanics and 1 in GTR

$$\begin{cases} \delta\dot{\Omega}_{\text{obs}}^{\text{Mercury}} = \dot{\Omega}_{.2}^{\text{Mercury}}\delta J_{2\odot} + \dot{\Omega}_{.4}^{\text{Mercury}}\delta J_{4\odot} + \dot{\Omega}_{LT}^{\text{Mercury}}\mu_{LT} + \Delta^{\text{Mercury}}, \\ \delta\dot{\Omega}_{\text{obs}}^{\text{Venus}} = \dot{\Omega}_{.2}^{\text{Venus}}\delta J_{2\odot} + \dot{\Omega}_{.4}^{\text{Venus}}\delta J_{4\odot} + \dot{\Omega}_{LT}^{\text{Venus}}\mu_{LT} + \Delta^{\text{Venus}}, \\ \delta\dot{\Omega}_{\text{obs}}^{\text{Mars}} = \dot{\Omega}_{.2}^{\text{Mars}}\delta J_{2\odot} + \dot{\Omega}_{.4}^{\text{Mars}}\delta J_{4\odot} + \dot{\Omega}_{LT}^{\text{Mars}}\mu_{LT} + \Delta^{\text{Mars}}. \end{cases} \quad (27)$$

The coefficients $\dot{\Omega}_{\ell}$ for the first two even zonal harmonics are

$$\begin{cases} \dot{\Omega}_{.2} = -\frac{3}{2}n\left(\frac{R_{\odot}}{a}\right)^2 \frac{\cos i}{(1-e^2)^2}, \\ \dot{\Omega}_{.4} = \dot{\Omega}_{.2} \left[\frac{5}{8} \left(\frac{R}{a}\right)^2 \frac{(1+\frac{3}{2}e^2)}{(1-e^2)^2} (7\sin^2 i - 4) \right] \end{cases} \quad (28)$$

$R_{\odot}$ is the Sun's equatorial radius. The quantities Δ in eq.(27) refer to the other unmodelled or mismodelled effect which affect the temporal evolution of the nodes of the considered planets. In the present case they would be the N -body gravitational interactions—of Newtonian origin—with the other planets and asteroids of the Solar System. From the nominal values of $\dot{\Omega}_{N\text{-body}}$ quoted in Table 3 it can be noted that their secular components are very important perturbing effects which should be known with very high accuracy in order to prevent a fatal aliasing effect on the proposed measurement.

If we solve eq.(27) for the Lense-Thirring parameter μ_{LT} it is possible to obtain

$$\begin{cases} \delta\dot{\Omega}_{\text{obs}}^{\text{Mercury}} + k_1\delta\dot{\Omega}_{\text{obs}}^{\text{Venus}} + k_2\delta\dot{\Omega}_{\text{obs}}^{\text{Mars}} = X_{LT}\mu_{LT} + (\text{other secular mismodelled effects } \Delta), \\ k_1 = \frac{\dot{\Omega}_{.2}^{\text{Mars}}\dot{\Omega}_{.4}^{\text{Mercury}} - \dot{\Omega}_{.2}^{\text{Mercury}}\dot{\Omega}_{.4}^{\text{Mars}}}{\dot{\Omega}_{.2}^{\text{Venus}}\dot{\Omega}_{.4}^{\text{Mars}} - \dot{\Omega}_{.2}^{\text{Mars}}\dot{\Omega}_{.4}^{\text{Venus}}} = -48.008308, \\ k_2 = \frac{\dot{\Omega}_{.2}^{\text{Mercury}}\dot{\Omega}_{.4}^{\text{Venus}} - \dot{\Omega}_{.2}^{\text{Venus}}\dot{\Omega}_{.4}^{\text{Mercury}}}{\dot{\Omega}_{.2}^{\text{Venus}}\dot{\Omega}_{.4}^{\text{Mars}} - \dot{\Omega}_{.2}^{\text{Mars}}\dot{\Omega}_{.4}^{\text{Venus}}} = 510.404066, \\ X_{LT} = \dot{\Omega}_{LT}^{\text{Mercury}} + k_1\dot{\Omega}_{LT}^{\text{Venus}} + k_2\dot{\Omega}_{LT}^{\text{Mars}} = 0.002069'' \text{ cy}^{-1}, \end{cases} \quad (29)$$

where the numerical values of the coefficients k_1 and k_2 and the slope X_{LT} of the gravito-magnetic trend come from the values of Table 3. The meaning of eq.(29) is the following. Let us construct the time series of the residuals of the nodes of Mercury, Venus and Mars by using real observational data and the full dynamical models in which GTR is purposely set equal to zero, e.g. by using a very large value of c . We expect that, over a multidecadal observational time span, the so combined residuals will fully show the GTR signature and partly the mismodelled N -body effect¹³ in terms of a linear trend. The measured slope, divided by X_{LT} , yields μ_{LT} which should be equal to one if GTR was correct and if the bias from the residual N -body effect was sufficiently small. If we use the results of Table 4 in eq.(29) in order to get an idea of what could be the observational accuracy of such a measurement we get a 19% error. It is important to note that such an estimate does not include the systematic error induced by mismodelled secular effect due to the classical N -body interactions.

6.2.2. A $J_{2\odot} - (N\text{-body})$ free combination

Eq.(29) is designed in order to cancel out the contributions of the first two even zonal harmonics of the multipolar expansion of the Sun's gravitational potential. However, it is affected by the N -body classical secular precessions which, as it turns out from the figures in Table 3, are quite large; their mismodelled part, which would be accounted for by the orbital residuals, could be turn out to be very insidious with respect to our proposed measurement of the Lense-Thirring effect.

In view of these considerations, it would be better to set up another observable which is insensitive just to such huge classical secular effects. Moreover, from the results of Table 3 and from the evaluations of (Rozelot et al. 2004) according to which the possible magnitude of $J_{4\odot}$ would span the range $10^{-7} - 10^{-9}$, the secular precession induced by the octupolar mass moment of the Sun is negligible with respect the Lense-Thirring rates. These facts lead us to

¹³ Possible aliasing time-dependent N -body residual effects with the periodicities of the outer planets, mainly Jupiter, should average out over a sufficiently long multidecadal time span. However, it would be possible to fit and remove them from the time series.

design a three-node combination which cancels out the effects of $J_{2\odot}$ and of the classical N -body precessions and is affected by $J_{4\odot}$. It can be obtained by substituting $\dot{\Omega}_{.4}$ with $\dot{\Omega}_{\text{class}}$ in eq.(29). It is

$$\begin{cases} \delta\dot{\Omega}_{\text{obs}}^{\text{Mercury}} + k'_1\delta\dot{\Omega}_{\text{obs}}^{\text{Venus}} + k'_2\delta\dot{\Omega}_{\text{obs}}^{\text{Mars}} \sim Y_{\text{LT}}\mu_{\text{LT}}, \\ k'_1 = \frac{\dot{\Omega}_{.2}^{\text{Mars}}\dot{\Omega}_{\text{class}}^{\text{Mercury}} - \dot{\Omega}_{.2}^{\text{Mercury}}\dot{\Omega}_{\text{class}}^{\text{Mars}}}{\dot{\Omega}_{.2}^{\text{Venus}}\dot{\Omega}_{\text{class}}^{\text{Mars}} - \dot{\Omega}_{.2}^{\text{Mars}}\dot{\Omega}_{\text{class}}^{\text{Venus}}} = -10.441702, \\ k'_2 = \frac{\dot{\Omega}_{.2}^{\text{Mercury}}\dot{\Omega}_{\text{class}}^{\text{Venus}} - \dot{\Omega}_{.2}^{\text{Venus}}\dot{\Omega}_{\text{class}}^{\text{Mercury}}}{\dot{\Omega}_{.2}^{\text{Venus}}\dot{\Omega}_{\text{class}}^{\text{Mars}} - \dot{\Omega}_{.2}^{\text{Mars}}\dot{\Omega}_{\text{class}}^{\text{Venus}}} = 9.765758, \\ Y_{\text{LT}} = \dot{\Omega}_{\text{LT}}^{\text{Mercury}} + k'_1\dot{\Omega}_{\text{LT}}^{\text{Venus}} + k'_2\dot{\Omega}_{\text{LT}}^{\text{Mars}} = -0.000351'' \text{ cy}^{-1}, \end{cases} \quad (30)$$

The systematic error affecting eq.(30) should be totally negligible because it would be due only to the higher degree multipole mass moments of the Sun. The observational error, according to the results of Table 4, would amount to 36%.

7. Conclusions

In this chapter we have proposed a relatively cheap approach in testing post-Newtonian gravity in the Solar System scenario with particular emphasis on some secular gravito-electromagnetic effects on the orbital motion of the planets. It is based on a reanalysis and a suitable combination of the ranging data to planets. As a by-product, also the important solar quadrupole mass moment $J_{2\odot}$ could be accurately measured.

First, we have explicitly worked out the post-Newtonian gravito-electric secular rate of the mean anomaly of a test particle freely orbiting a spherically symmetric central object. Moreover, we have outlined a possible strategy for determining simultaneously and independently of each other, the solar quadrupole mass moment $J_{2\odot}$ and two parameters ν_{GE} and μ_{GE} which account for the post-Newtonian gravito-electric secular shifts of the perihelion and the mean anomaly, respectively, of planets. They are 0 in Newtonian mechanics and 1 in the General Theory of Relativity. They could be expressed in terms of the standard PPN γ and β parameters which could, then, be determined independently of each other as well. The usual approach employed in the ephemerides data reductions tests post-Newtonian gravity theories as a whole straightforwardly in terms of the PPN parameters involving the simultaneous fit of many more or less correlated astrodynamical parameters among which there are also γ , β and $J_{2\odot}$. In this case, instead, we propose to analyze the time series of three suitably linearly combined residuals of $\dot{\Omega}$, $\dot{\pi}$ and $\dot{M}$ of Mercury, Venus and Mars built up in order to single out just certain selected Newtonian and post-Newtonian orbital effects. This can be done by setting purposely equal to zero (or to some default values to be subsequently adjusted with the proposed strategy) the orbital effects of interest in the force models of the equations of motion. By suitably choosing the length of the orbital arcs it would be possible to account for the secular terms only. The coefficients of the combinations would make each of them sensitive just to one orbital effect, independently of the other ones. By fitting the experimental residual signals with straight lines, measuring their slopes in $'' \text{ cy}^{-1}$ and suitably normalizing them would yield the values of $J_{2\odot}$, μ_{GE} and ν_{GE} . The obtainable formal accuracy would be of the order of $10^{-4} - 10^{-5}$ for γ and β and, more interestingly, 10^{-9} for $J_{2\odot}$.

Second, we have extended this strategy to the gravito-magnetic Lense-Thirring effect induced by the Sun's angular momentum. We have designed two linear combinations which would allow to measure such tiny effect, by construction, independently of the aliasing impact of the Sun's oblateness and of the N -body precessions. The obtainable accuracy would be of the order of 20-30%.

However, it must be pointed out that many of the proposed tests involve the analysis of the orbital motion of Mercury. The quality and the accuracy of its ephemerides should increase in future thanks to the already launched Messenger and planned BepiColombo spacecrafts, so that it is reasonably presumable that the precision of the proposed tests will increase as well.

Acknowledgements. L. Iorio is grateful to L. Guerriero for his support while at Bari. Special thanks to E. M. Standish and E. Pitjeva for their help and useful discussions and clarifications, and to W.-T. Ni for the updated reference about ASTROD.

References

- Anderson, J. D., Jurgens, R. F., Lau, E. L., & Slade, M. A. 1996, *Icarus*, 124, 690
- Armstrong, J., & Kuhn, J. R. 1999, *ApJ*, 525, 533
- Bertotti, B., Iess, L., & Tortora P. 2003, *Nature*, 425, 374
- Ciufolini, I. 1986, *Phys. Rev. Lett.* 56, 278
- Ciufolini, I. 1991, *Int. J. of Mod. Phys. A*, 6, 5511

- Ciufolini, I., & Wheeler, J. A. 1995, *Gravitation and Inertia* (Princeton University Press, New York)
- Ciufolini, I. 1996, *Il Nuovo Cimento A*, 109, 1709
- Ciufolini, I., Pavlis, E. C., Chieppa, F., Fernandes-Vieira, E., & Pérez-Mercader, J. 1998, *Science*, 279, 2100
- De Sitter, W. 1916, *MNRAS*, 76, 699
- Einstein, A. 1915, *Preuss. Akad. Wiss. Berlin Sitzber.*, 47, 831
- Elsworth, Y., Howe, R., Isaak, G. R., McLeod, C. P., Miller, B. A., New, R., Wheeler, S. J., & Gough, D. O. 1995, *Nature*, 376, 669
- Everitt, C. F. W., Buchman, S., DeBra, D. B., and other members of the Gravity Probe B Team 2001, *Gravity Probe B: Countdown to Launch. In Gyros, Clocks, and Interferometers: Testing Relativistic Gravity in Space*, ed. C. Lämmerzahl, C. W. F. Everitt, & F.W. Hehl, (Springer-Verlag, Berlin), 52
- Iorio, L. 2002, *Class. Quantum Grav.* 19, 5473
- Iorio, L. 2003, *Celest. Mech & Dyn. Astron.* 86, 277
- Iorio, L., Lucchesi, D.M., & Ciufolini, I. 2002a, *Class. Quantum Grav.* 19, 4311
- Iorio, L., Ciufolini, I., & Pavlis, E. C. 2002b, *Class. Quantum Grav.*, 19, 4301
- Iorio, L., & Morea, A. 2004, *Gen. Rel. Gravit.* 36, 1321
- Iorio, L., Ciufolini, I., Pavlis, E.C., Schiller, S., Dittus, H., & Lämmerzahl, C. 2004, *Class. Quantum Grav.* 21, 2139
- Jacobson, R. A. 2003, JUP230 orbit solution
- Lense, J., & Thirring, H. 1918, *Phys. Z.* 19, 156 translated by Mashhoon, B., Hehl, F. W., & Theiss, D. S. 1984, *Gen. Rel. Grav.*, 16, 711
- Lucchesi, D. M. 2003, *Phys. Lett. A*, 318, 234
- Mashhoon, B., Iorio, L., & Lichtenegger, H. 2001, *Phys. Lett. A*, 292, 49
- Melliti, T., Fridelance, F., & Samain, E. 2002, *A&A*, at press
- Milani, A., Vokrouhlický, D., Villani, D., Bonanno, C., & Rossi, A. 2002, *Phys. Rev. D*, 66, 082001
- Ni, W.-T., Shiomi, S., & Liao, A.-C. 2004, *Class. Quantum Grav.*, 21, S641
- Nordvedt, K. 2001, *An Overview of Solar System Gravitational Physics: The Theory-Experiment Interface*, In *Gyros, Clocks, and Interferometers: Testing Relativistic Gravity in Space*, ed. C. Lämmerzahl, C. W. F. Everitt, & F.W. Hehl, (Springer-Verlag, Berlin), 3
- Nordvedt, K. 2003, preprint gr-qc/0301024
- Paternó, S., Sofia, S., & Di Mauro, M.P. 1996, *A&A*, 314, 940
- Pettengill, G. H., Eliason, E., Ford, P. G., Lorient, G. B., Masursky, H., & McGill, G. E. 1980, *J. Geophys. Res.*, 85, A13, 8261
- Pijpers, F. P. 2003, *A&A*, 402, 683
- Pireaux, S., & Rozelot, J.-P. 2003, *Ap&SS*, 284, 1159
- Pitjeva, E. V. 2001a, *Cel. Mech. & Dyn. Astron.*, 80, 249
- Pitjeva, E. V. 2001b, *A&A*, 371, 760
- Ries, J. C., Eanes, R. J., & Tapley, B. D. 2003, *Lense-Thirring Precession Determination from Laser Ranging to Artificial Satellites. In Nonlinear Gravitodynamics. The Lense-Thirring Effect*, ed. R.J. Ruffini, & C. Sigismondi, (World Scientific, Singapore), 201
- Rozelet, J.-P., Pireaux, S., Lefebvre, S., & Corbard, T. 2004, preprint astro-ph/0403382
- Schiff, L.I. 1960 *Am. J. Phys.*, 28, 340
- Shapiro, I., Pettengill, G. H., Ash, M. E., Ingalls, R. P., Campbell, D. B., & Dyce, R. B. 1972, *Phys Rev. Lett.*, 28, 1594
- Shapiro, I., Counselman, C., & King, R. 1976, *Phys Rev. Lett.*, 36, 555
- Shapiro, I. 1990, *Solar system tests of general relativity: recent results and present plans. In Proceedings of the 12th International Conference on General Relativity and Gravitation*, ed. N. Ashby, D. Bartlett, & W. Wyss, (Cambridge University Press, Cambridge), 313
- Soffel, M. H. 1989, *Relativity in Astrometry, Celestial Mechanics and Geodesy* (Springer Verlag, Heidelberg)
- Standish, E. M. 1998, *JPL planetary and lunar ephemerides DE405/LE405*, Interoffice Memorandum 312.F-98-048, 1
- Standish, E. M. 2002, *Future JPL Ephemerides. In Proceedings of the IERS Workshop on the Implementation of the New IAU Resolutions*, IERS Technical Note No. 29, ed. N. Capitaine, D. Gambis, D. D. McCarthy, G. Petit, J. Ray, B. Richter, M. Rothacher, E. M. Standish, & J. Vondrak, (Verlag des Bundesamts für Kartographie und Geodäsie, Frankfurt am Main), 61
- Stairs, I. H., Arzoumanian, Z., Camilo, F., Lyne, A. G., Nice, D. J., Taylor, J. H., Thorsett, S. E., & Wolszcan, A. 1998, *ApJ*, 505, 352
- Turyshev, S., Shao, M., & Nordvedt, K. 2004, *Class. Quantum Grav.*, 21, 2773
- Turyshev, S., Williams, J. G., Nordvedt, K., Shao, M., & Murphy, T. W. 2003, *35 Years of Testing Relativistic Gravity: Where do we go from here? invited talk presented at 302.WE-Heraeus-Seminar: "Astrophysics, Clocks and Fundamental Constants"*, Bad Honnef, Germany, June 16-18, 2003, preprint gr-qc/0311039
- Vecchiato, A., et al. 2003, *A&A*, 399, 337
- Will, C. M. 1993, *Theory and Experiment in Gravitational Physics*, 2nd edition (Cambridge University Press, Cambridge)
- Xu, X., & Ni, W.-T. 1997, *Solar angular momentum calculation. In National Science Council report*